

In

Dysentery

Inaugural Dissertation

on the

Dysentery

by

Archibald D. Moultrie

of Virginia

April the 10th 1841



P

ance about
of autumn
remittent
however oc
country, a
mies, to g
At genes
inability
appetite, a
mouth, we
toms are
regular ac

UNIVERSITY
PENNSYLVANIA
LIBRARY

Dysentery

Generally makes its appearance about the end of summer, or beginning of autumn, at which time intermittent, and remittent fevers, are most prevalent. — It may however occur at any time, or season, or in any country, and is subject, like all other epidemics, to great variety in appearance and cure.

It generally begins with great lassitude, and inability to motion, chilliness and thirst, loss of appetite, and sometimes a bitter taste in the mouth, with nausea and vomiting. — These symptoms are very soon succeeded, by morbid or irregular action in the arterial system, attended

with more
and believe
At sometime
of affection
-ness, and
those arising
-asionally
from the
-ted, gener
In some co
ul any d
-tee of by
dry dysent
Rush, who
so great in
as to render

with severe pain in the head, excessive thirst,
and bilious stools.—

It sometimes however, appears with a variety
of affections in the alimentary canal, as costi-
venae, and not unfrequently pains resembling
those arising from colic. And a purging is oc-
casionally the first symptom, but this is not
often the case, for the symptoms above enumera-
ted, generally usher in the disease.—

In some cases severe dysenteric pains come on, with-
out any discharge whatever. This was taken no-
tice of by Doctor Sydenham, and called by him
dry dysentery— and has been accounted for by Dr
Rush, who supposes that the force of the disease is
so great in the bowels, as to prostrate them, so much,
as to render them incapable of acting on their contents,

until the vaci-
ing or purging
observes a w
efs of morbi
kidneys, in
a constituty
By neglecting
-lly gains
the patient,
griping, and
ing this, no
ged with
short respite
The matter
sembling a
blood and

until the excitement is lessened by means of bleed-
ing or purging. - It is not in dysentery alone we
observe a want of secretion or excretion from ex-
cess of morbid excitement. - It takes place in the
kidneys, in the liver, in the eyes, and in gonorrha-
a, constituting what is called a dry clap. -

By neglecting the above symptoms, the disease gradu-
ally gains strength, without exciting much fear in
the patient, until he is attacked with flatulence,
gripping, and inclination to go to stool; in indulg-
ing this, nothing is voided, but a little mucus ten-
ged with blood; after this there is generally a
short respite from pain, but this ^{is} of short duration.

The matter voided by stool is various; at first re-
sembling a purging of thin excrement, sometimes
blood and mucus, mucus alone, blood and faeces,

and in some
and not for
cylindrical au
elling suit,
created, wh
There is co
Early the
resp, and n
upper part
the gripping
the firm w
considerable
his disease,
lows any
instant in
had taken

and in some cases blood alone, this comes from ~~enlarged~~
and not from ruptured vessels, as has been supposed,
scybala are sometimes discharged, also balls resum-
bling suet, a membrane is not unfrequently eva-
cuated, which is the effect of inflammation. —
There is considerable pain in the bowels particu-
larly the colon and rectum, and great uneasi-
ness, and nausea at stomach, which shews that the
upper part of the intestinal canal is affected. —
The griping and tenesmus continuing to increase,
the fever which is of the remitting kind, becomes more
considerable. — A symptom has been remarked in
this disease, that is to say, when the patient swal-
lows any thing either solid, or liquid, he has an
instant inclination to go to stool, as if what he
had taken in, passed immediately through him.

but it will
usual disor
had been su
stability of
velop some
every sympt
frequent an
pined wi
Nothing se
cathartic s
an evacuat
the expuls
the distor
long, for the
nolered a
the colera

but it will be found upon examination, to be the usual discharge attending the disease, and not what had been swallowed; this is owing to the great irritability of the intestinal canal. —

Unless something be done for the patient at this time, every symptom increasing, the stools will become more frequent and painful, with a fetid smell, accompanied with severe griping, and tenesmus. —

Nothing accrementitious is discharged, unless a cathartic be administered, when small, round balls are evacuated, termed by Authors scybala. — Upon the expulsion of these, there is an alleviation of the distressing symptoms, but this does not continue long, for the complaint increasing with redoubled violence a sebaceous matter is discharged, resembling the liliæ carniū, or washings of flesh, with an

extremely fat
The pulse
and tense, n
countenance
their accuston
during sleep,
covered with
or, with great
The debility
of weak, frequ
the evacuation
continues to
ing and done
great, sudden
and flattered
The delusion

extremely fetid and cadaverous smell.—

The pulse which until this time, was hard, full, and tense, now becomes small and frequent, the countenance appears dejected, and the eyes lose their accustomed lustre, and appear only half-closed during sleep, the tongue and teeth become foul and covered with a dark, tenacious scurf, hiccough comes on, with great prostration of strength.—

The debility progressing, the pulse becomes exceedingly weak, frequent and almost imperceptible.—

The evacuations are often involuntary, and the smell continues to be extremely offensive.— The pain, griping and tenesmus which until now had been very great, suddenly ceasing, the patient and his friends are flattered with the hopes of a speedy recovery.— The delusion however exists but for a short time,

for the nau

become cold

if not attog

comes on, ap

schremities

and the pa

The appeara

times meet

world action

secondly, mar

fourthly, just

somewhat, the

eightly, the la

spleen is said

The remote

those producing

for the nausea and vomiting returning, the sweats become cold and clammy, deglutition difficult, if not altogether impossible, subcutis tendinum comes on, aphthae make their appearance, the extremities become cold, delirium supervenes, and the patient dies.

The appearances on dissection, are various, we sometimes meet with no marks of disease whatever, from morbid action transcending the grade of inflammation, secondly, marks of inflammation, thirdly, a membrane, fourthly, pustules, fifthly, tubercles, sixthly, gangrene, seventhly, the inner coat of the intestine abraded, eighthly, the liver found generally diseased, ninthly, the spleen is said to be enlarged.

The remote causes of dysentery are the same with those producing bilious and yellow fever, and is in fact,

only a modification
furnished with
prove, that is
quently term
Dr. Blegborn
suffered, b
mon for dys
tians, and f
regularly atte
occurring abo
rally; proves
contagion, b
and yellow fe
reasons, that
are to be ma
The predispo

only a modification of the same disease. For we are furnished with many instances, which incontestably prove, that intermittent and remittent fevers frequently terminate in dysentery, and vice versa. Dr. Bleghorn tells us, when one of these diseases is suppressed, the other often ensues; nor is it uncommon for dysenteric fevers to put on the form of tertians, and for the paroxysms of tertians to be regularly attended with gripes and stools. For its occurring about the same season of the year generally, proves that it does not depend upon specific contagion, but an impure atmosphere, like bilious, and yellow fevers. For these and innumerable other reasons, that could be given, I consider the remote causes to be marsh and human effluvia.

The predisposing cause of dysentery, is, debility, whether

induced directly
the depressing
despair - which
is a powerful
illness - the
fruitful source
of error in
in support of
has no doubt
the disease
and may pro-
ceeds to pro-
duce of people
the common
affected, both
than those who

induced directly or indirectly. - The direct causes, are the depressing passions of the mind, as fear, grief and despair - Cold. This is universally acknowledged to be a powerful cause in inducing debility.

Filthiness. This has always been considered as a fruitful source of this disease, and its frequent occurrence in camps and hospitals, has been adduced in support of this opinion. - A neglect of cleanliness, has no doubt, a considerable share in rendering the disease more violent, when it has taken place, and may probably sometimes concur with other causes to produce it; for we find that the poorer class of people, many of whom, are destitute of the common necessaries of life, become much sooner affected, both with this and many other diseases, than those who enjoy a higher station. -

The intestines
frequency of
weather. - 2
8th unusual
at more in
ability in
Improper a
of the food
are more s
in circulation
live in a C
gestion, and
negotiable ac
In a debilit
the intestines
to create the
from their ^{living} ^{up}

The indirect causes, are 1.st Heat, hence the greater frequency of dysentery in warm climates, and warm weather. - 2.nd Intemperance in eating and drinking. 3.rd Unusual labour or excess. - The causes which act more immediately in inducing a state of debility in the stomach and intestines, are 1.st Improper aliment, as the immoderate use of unripe fruit. - This is ^{the} cause, why country people are more subject to this disease, than those who live in a city. ^{an exhalation from the putrefaction} 2.nd Animal food of difficult digestion, and of an unwholesome quality. 3.rd The vegetable acids. 4.th Unwholesome water. -

In a debilitated, and consequently excitable state of the intestines, it only requires the action of stimuli to create the disease in question, and this happens from their ^{being} either immediately, or more remotely applied.

It has been
vicarious respiration
It is unnecessary
which have
appears to have
the disease,
in the lungs
most satisfac-
the perimate
ly of the in-
in an irreg-
system, but
of the weak
and hives, &
hence they
operations,

There has been great diversity of opinion among physicians respecting the proximate cause of dysentery. It is unnecessary to mention the various theories which have been advanced. — But Dr Sydenham appears to have been the first who had a correct idea of this disease, he considered it as a fever turned in upon the bowels. — But as Dr Rush has given us the most satisfactory theory, I shall mention his, namely, the proximate cause of fever in general, and consequently of the intestinal state of fever, appears to consist in an irregular, convulsive action of the arterial system, but this irregular action, in consequence of the weak and debilitated state of the stomach, and bowels, is particularly determined upon them, hence they suffer from pain, inflammation, sup-
puration, gangrene &c.

And is perhaps

it can be shown

under considera-

more certainly

and when given

In dysentery

various, or suite

But in the tr

to be of a sym

case, the follo

Bloodstiffing

or when the

indispensable

tinued, until

try symptoms

do. - In deter

There is perhaps no disease in which the aid of a physician is more essentially requisite, than in the one under consideration, for there is no disease which proves more certainly fatal, when left in the hands of nature; and when judiciously managed none more easily cured.

In dysentery as in most diseases our practice must be various, & suited to the different states of the system;

But in the United States dysentery is generally found to be of a very inflammatory nature, when this is the case, the following mode of practice will be found best.

Bloodletting when the pulse is full, quick and tense, or when the patient complains of great pain, is indispensably necessary, the bleeding should be continued, until the pulse is subdued or the inflammatory symptoms have abated, and the pain be relieved. In determining the quantity of blood to be drawn

no position
upon the
the habit.
Emetics
show the
really wh
of bile, fo
be found, t
Purgatives
upon three
in a great
post, by
and second
from the
new and
greatest

no positive rule can be established, it must depend upon the judgment of the physician, together with the habit, age and constitution of the patient.

Emetics are seldom useful, tho they may be used where the patient complains of nausea, and especially when this depends upon a redundancy of bile, for this purpose Speccacanthus perhaps will be found best.

Purgatives have been justly recommended, and upon their judicious administration, the cure will in a great measure depend, they act in two ways, first, by reducing the excitement in the bloodvessels, and secondly by discharging the irritating matter from the intestines which is the cause of the tenesmus and griping—Clarke and Pringle place the greatest reliance on purges, and even advise their

Early we,
removed, the
and the n
ended Con
or bowls—
and is cer
however we
has been a
may be en
of this meo
by Doctor C
thy form n
Having po
medies pe
or a powder
this powder

daily use, until the most troublesome symptoms are removed; they should be of castor, or sweet oil, sulphur, and the neutral salts. Sulphur has been recommended (on the supposition of its acting on the lower bowels. Calomel has been highly spoken of, and is certainly entitled to our attention, of this however we shall speak hereafter. — Rheubarb has been used; and under certain circumstances may be employed with advantage, an infusion of this medicine has been said to act as a tonic, by Doctor Barton, and after sufficient depletion this form may perhaps be used with advantage. Having promised sufficient evacuations, various remedies present themselves, such as the chalk pulv., or a powder composed of opium, Specac, and creta; this powder has been highly spoken of by D^r Barton

in bloody
Tudor's
try diath
as a dis
force, and
James p
operation
by water
Injections
they shou
with the
relieves th
the imitat
the operat
Opium. in
company, it

in bloody dysentery.

Sudorifics may be given when the inflammatory diathesis is subdued, or when the system shows a disposition to throw off the disease by the pores, and the best sudorifics will be Dovers or James's powders, they should be assisted in their operation by diluents, such as wine whey, barley water, or flaxseed tea.

Injections may be used in every stage of the disease, they should be of an oily or mucilaginous nature with the addition of a little laudanum, which relieves the griping and tenesmus, they bring away the irritating matter from the intestines, and assist the operation of purgatives.

Opium in the cure of dysentery is indispensably necessary, it should be given in small doses during the

any and lo
Sleep, this
meager
History, in
essential se
the skin an
to the wis
Delicents a
He commu
it were, the
their cont
Finally, wh
should be s
attention
by being
part of

day and large ones at night to procure ease and sleep, this it does more effectually than any other remedy.

Plasters, in this as in many other diseases, are of essential service; they act by easing pain, relaxing the skin and inducing sleep, they may be applied to the wrists, ancles, thighs or abdomen.

Delicents and demulcents should be given from the commencement of the disease; they sheath, as it were, the intestines, and blunt the acrimony of their contents.

Finally, when the above remedies fail, mercury should be resorted ^{to}, so as to produce a gentle salivation; the disease, is, in this way cured, by being translated from the bowels, to a part less essential to life.

[Faint, illegible handwriting, likely bleed-through from the reverse side of the page.]

C

11

The

The

For the

On the

C

of